

**Instructions for Fish Antibiotics in Humans**  
**PensacolaPreppers**

**Ampicillin**

Urinary Tract Infections

Adult Dosage: 500 milligrams, 4 times a day. Duration: 3 days.

**Amoxicillin**

Ear Infections

Bronchitis (treat and if no improvement seen change to Keflex)

Strep Throat (treat and if no improvement seen change to Keflex)

Skin Infections (if not bad)

Adult Dosage: 500 milligrams, 2 times a day. Duration: 7 days.

**Keflex**

**Note:** This is much stronger than amoxicillin and can result in severe diarrhea.

Wherever possible, use amoxicillin first. Use Keflex as last resort!

Bronchitis

Stomach bugs (diarrhea and/or vomiting)

Ear Infections

Adult Dosage: 500 milligrams, 2 to 4 times a day, with food. Duration: 3 to 7 days.

**Metronidazole**

For reactions to antibiotics, especially Keflex.

If taking antibiotics and you get severe diarrhea then take Metronidazole with the anti-biotic to counter the bad side affects of the anti-biotic.

Some Vaginal problems (bacterial)

Adult Dosage: 250 milligrams 4 times a day. Duration: 7-14 days  
Do not drink alcohol.

**Penicillin:**

Mild Infections, skin, cuts, gun shots

Note: If redness creeps past mark (grows), switch to Keflex or something else.

Adult Dosage: 500 milligrams, 4 times per day. Duration: 7 – 10 days.  
One hour before meal or two hours after meal.